



Evaluation of Supplementation of *Embelica officinalis* on Growth Parameter, Yield and Biological Efficiency of *Pleurotus sajor-caju*.

Poonam Dehariya and Deepak Vyas

Lab of Mushroom Biology, Department of Botany,
Dr. H. S. Gour Vishwavidyalaya, Sagar - 470 003, Madhya Pradesh, India

Received: 05 May 2020; Revised accepted: 15 August 2020

ABSTRACT

Six substrates viz. soybean straw, wheat straw, paddy straw, domestic waste, fruit waste and used tea leaves was taken for the purpose. For the improvement of yield supplementation of *Embelica officinalis* was done. Among all test substrates soybean straw was give maximum yield 935.0 gm/kg, 956.7 gm/kg, 96.0 gm/kg at 1%, 2% and 3% supplementation respectively. It was observed from the data that different concentrations of *Embelica officinalis* was proved good for the enhancement of yield but 3% concentration was given maximum yield and biological efficiency.

Key words: Supplementation, Yield, Biological efficiency, *Embelica officinalis*

Oyster mushroom is one of the most popular edible mushroom and belong to the genus *Pleurotus* and the family *Pleurotaceae* (Flack 1917) and is now grown commercially around the world for food. Oyster mushroom is one of the most commonly sought wild mushrooms, though it can also be cultivated on straw and other media. *Pleurotus* mushrooms are primary decomposers of hardwood trees and are found worldwide. Fruiting bodies as well as active mycelia of *Pleurotus* species also possesses a number of therapeutic properties like anti-inflammatory, immunostimulant and anticancer activity, immunomodulatory, ribonuclease activity and many other activities (Patel *et al.* 2012). *P. sajor-caju* is one of the most successfully cultivated species of these mushrooms and it is considered to be delicious (Zhang *et al.* 2002). Agriculture residues are the major source of lignocellulosic materials, which is best substrate for solid state fermentation of edible fungi such as *P. sajor-caju*. Zadrazil (1980) showed that *P. sajor-caju* has very high saprophytic colonizing ability and can degrade wheat straw efficiently. Bisaria *et al.* (1987) studied the growth of this mushroom on several different agricultural wastes, including paddy and wheat straws. Growing oyster mushrooms convert a high percentage of the lignocellulosic

materials to fruiting bodies. Therefore, cultivation of *P. sajor-caju* on various agricultural residues offers high value products with nutritional and medicinal properties (Pokhrel *et al.* 2013).

Mushroom supplementation is an agronomic process which consists of the application of nutritional amendments to the substrates employed for mushroom cultivation. Different nitrogen and carbohydrate rich supplements have been evaluated in crops with a substantial impact on mushroom yield and quality. The addition of external nutrients increases the productivity of some low-yielding mushroom varieties, and therefore is a useful tool for the industry to introduce new commercially viable varieties. On the other hand, a new line of research based on the use of mushroom growth promoting microorganisms is rising above the horizon to supplement the native microbiota, which appears to cover nutritional deficiencies. Several supplements employed for the cultivated mushrooms and their agronomic potential in terms of yield and quality of the crop and to design new formulas for commercial supplementation (Carrasco *et al.* 2018). Supplementation has been reported to cause a rise in substrate temperature, possibly due to faster metabolic activities triggered by extra nitrogen. Royse and Schisler (1986) observed overheating (from 30 to 47°C) in bags where a nitrogen-rich supplement was applied without benomyl (fungicide) treatment and proposed that it could be due to the growth of competitor moulds.

***Corresponding author:** Dr. Poonam Dehariya, Assistant Professor, Department of Botany, Dr. H. S. Gour Vishwavidyalaya, Sagar - 470 003, Madhya Pradesh

e-mail: dpoonambot18@gmail.com

Addition of the supplements with basal substrate has been as common practice to enhance the yield, nutritional and medicinal values. In oyster mushroom cultivation, multiple basal substrates and supplements (additives) are reported to be used by researchers. Therefore, the technology of mushroom agriculture enables us to acquire substrate materials at very low cost or even for free to get biotechnological foods and further led to conserve our environment through efficient bioconversion of wastes with sustainable food security (Naraian *et al.* 2016). Present study was therefore conducted to determine the effects of different substrates viz. Soybean straw, wheat straw, paddy straw, domestic waste, fruit waste and used tea leaves and *Embelica officinalis* supplements on biological efficiency and yield parameters of *Pleurotus sajor caju*.

MATERIALS AND METHODS

Spawn preparation

Clean whole grains were taken for the purpose. The grains were pre-wetted by boiling in water for 20-30 min (Jain 2005). After boiling, excess water was drained off by spreading the grains on a wire mesh. Grains were now mixed with gypsum (calcium sulphate) and chalk powder (calcium carbonate) at the rate of 2% and 0.5%, respectively on dry weight basis. The grains were filled in flasks, and plugged then sterilized in autoclave at 22 Lb pressure for 1.5-2 hours. The grains were allowed to cool in room temperature for overnight. Next day flasks were inoculated with two bits of agar medium colonized with the mycelium of pure cultures. About 7-10 days after inoculation, flasks were shaken vigorously. Three weeks after incubation, the stock culture becomes ready for further multiplication of spawn. Inoculated flasks were incubated at $26 \pm 2^\circ\text{C}$.

Cultivation

A medium was prepared using soybean straw (SS), wheat straw (WS), paddy straw (PS), domestic wastes (DW), Fruit waste (FW), used tea leaves (UTL). All substrates were washed in fresh water. The chopped straw substrates were steeped in water containing 75 ppm bavistin + 500 ppm formaldehyde for 18 hours (Jain 2005) for preventing mould infestation due to various other competing fungi. In the present investigation *Embelica officinalis* supplementation was used as 1, 2, 3 percent concentration on the dry weight basis of substrate respectively. Supplements were sterilized with 25 ppm bavistin for 12-18 hrs. Excess water was drained and was dried in shed to retain 65-70% moisture content by squeezing with hands and then allowed to cool down for a certain period (1 hr). Spawning was done @ 2% wet weight basis of substrate by thoroughly mixing. Spawned substrate was filled up in perforated polythene bags (60 × 40 cm) and polythene mouth was closed with rubber band. These bags were transferred to crop room for spawn run. Three replications were maintained for each substrate. For spawn run, temperature and relative humidity were maintained between 25-30°C and 65-90%, respectively. Polythene bags were cut open when the mycelium run was completed. The substrate beds were moistened by sprinkling of water thrice

a day which was stopped a day before harvesting. Average values of observation with respect to duration of spawn run, time taken for first harvest yield and total yield was recorded. Biological efficiency of mushroom on fresh weight basis was calculated by using formula given by Chang and Miles (1989).

$$\text{Biological efficiency (\%)} = \frac{\text{Yield of fruiting body (gm)}}{\text{Total weight of substrate used (gm)}} \times 100$$

RESULTS AND DISCUSSION

The data presented in (Table 1) shows the impact of supplementation of *Embelica officinalis* on the growth and yield of *P. sajor-caju* cultivated on three conventional substrates viz. soybean straw, wheat straw and paddy straw. It was observed that supplementation of 1% powder of *E. officinalis* improve growth parameter as well yield but significant change was recorded with 3% supplementation. As soybean straw was give 935.0 gm/kg, 956.7.0 gm/kg, and 960.0 gm/kg at 1%, 2% and 3% supplementation respectively. Paddy straw was give comparatively lower yield viz. 750.0 gm/kg, 761.7 gm/kg and 778.4 gm/kg at 1%, 2% and 3% supplementation respectively. The data presented in (Table 2) shows the impact of supplementation of *Embelica officinalis* on the growth and yield of *P. sajor-caju* cultivated on three non-conventional substrates viz. Domestic waste, Fruit waste and Used tea leaves. It was observed that among all the test three non-conventional substrates domestic waste was give maximum yield at 1%, 2% and 3% supplementation 730.0 gm/kg, 745.0 gm/kg, 763.4 gm/kg respectively. Used tea leaves shows comparatively lower yield which was 666.7 gm/kg, 671.7 gm/kg and 683.4 gm/kg respectively at 1%, 2% and 3% supplementation. Interesting trend of increasing yield was similar in conventional and non-conventional test substrates.

Addition of organic and inorganic supplements to other substrates are known to influence the yield of various species of mushroom (Jain 2005, Mishra and Singh 2007). The addition of protein-rich supplements is a common practice for nitrogen-deficient composts in the cultivation of mushrooms. Various researchers have used supplements from animal and plant origins, including protein-, carbohydrate- or oil-rich substances, for *Pleurotus* species (Gurjar and Doshi 1995).

Fungi contain direct absorption mechanism; thus, they utilize all the nutrients from plant material. Carbon and nitrogen plays an important role in the nutrition of fungi. Fungal species utilize various polysaccharides, monosaccharides, organic acids, amino acids, certain alcohols, polycyclic compounds, and natural products such as lignin and cellulose as carbon sources. The principle substrates for fungi in nature are plant materials. The plant cell wall consists of the polysaccharide's cellulose, lignin and hemicelluloses which are the carbon compounds utilized by fungi. These polysaccharides are insoluble and consequently they must be broken down to soluble units (monosaccharides) by the excretion of extracellular enzymes (e.g. cellulase, lignase) from the hyphae (Chang and Miles 2004).

Growth Parameter, Yield and Biological Efficiency of *Pleurotus sajor- caju*

Table 1 Effect of supplementation of different concentration of *Embelica officinalis* on growth parameter and yield of *P. sajor- caju* in selected conventional substrates

Substrates	Supplementation concentration (%)	Spawn run (days)	Pin head appearance (days)	Stipe length (cm)	Cap diameter (cm)	Total yield (gm/Kg)	BE (%)
Soybean straw	Control	17.66	22.4	2.9	7.8	933.4	93.3
	1%	17.0	21.4	2.9	7.8	935.0	93.5
	2%	16.7	20.7	3.0	7.9	956.7	95.6
	3%	16.0	20.0	3.1	8.0	960.0	96.0
Wheat straw	Control	19.0	23.4	2.6	8.1	800.0	80.0
	1%	18.0	22.7	2.7	8.7	810.0	81.0
	2%	18.16	21.7	2.8	8.8	821.7	82.1
	3%	17.0	21.4	2.9	9.0	850.0	85.0
Paddy straw	Control	21.4	25.4	2.7	8.0	743.4	74.3
	1%	21.0	24.7	2.8	8.1	750.0	75.0
	2%	20.0	24.4	2.9	8.3	761.7	76.1
	3%	19.0	23.0	2.9	8.4	778.4	77.8

Values are given in average of three replicates

Table 2 Effect of supplementation of different concentration of *Embelica officinalis* on growth parameter and yield of *P. sajor- caju* in selected non-conventional substrates

Substrates	Supplementation concentration (%)	Spawn run (days)	Pin head appearance (days)	Stipe length (cm)	Cap diameter (cm)	Total yield (gm/kg)	BE (%)
Domestic waste	Control	24.7	29.0	2.4	6.2	718.4	71.8
	1%	24.0	28.7	2.6	6.3	730.0	73.0
	2%	23.0	28.7	2.7	6.4	745.0	74.5
	3%	23.7	27.0	2.8	6.8	763.4	76.3
Fruit waste	Control	30.4	35.0	2.5	5.1	635.0	63.5
	1%	29.0	34.7	2.7	5.6	640.0	64.0
	2%	28.7	34.4	2.8	5.9	656.7	65.6
	3%	28.0	33.0	2.9	6.4	673.4	67.3
Used tea leaves	Control	25.7	31.7	2.3	5.6	655.0	65.5
	1%	25.0	31.0	2.4	5.7	666.7	66.6
	2%	24.7	30.7	2.5	5.9	671.7	67.1
	3%	24.0	30.0	2.6	6.1	683.4	68.3

Values are given in average of three replicates

Supplementation with various materials is recommended prior to spawning for enhancement of yield of Oyster mushrooms. Some of the recommendations are various oil seed meals and cakes, powdered pulses, wheat and rice bran etc. (Bahukhandi 1990), oil seed cake and meals, powdered pulses, sterilized chicken manure (Vijay and Upadhyay 1989, Baysal *et al.* 2003, Naraian *et al.* 2009), addition of oat meal (Jandaik 1974), rice bran (Gunasekharan and Graham, 1987), Wheat bran (Dies *et al.* 2003, Kalmis and Sargin 2004, Jeznabadi *et al.* 2016, Moonmoon *et al.* 2011) rice bran (Jednabadi *et al.* 2016, Peng *et al.* 2000, Vijay and Upadhyay 1989, Bonatti *et al.* 2003), oil seed cake (Shashirekh *et al.* 2002) broken corn and mineral solution (Moda *et al.* 2005), musterd seed cake (Naraian *et al.* 2009), soybean bran, soybean choker and groundnut bran (Dehariya *et al.* 2011), soybean choker (Jain and Vyas 2005), wheat bran and soybean flour (Jeznabadi *et al.* 2016), rice and wheat bran (Pedra and Marino 2006), cotton seed cake (Vijay and Upadhyay 1989), wheat bran and urea (Silva *et al.* 2007), Gupta and Raina (2008) reported effect of supplements on the yield of some high adaptive *Pleurotus species* in sub-tropics of Jammu. Perhaps the level of nitrogen content might be one of the

most important criteria to decide one of the positive effects on mushroom. The addition of amendments to the substrates influence the sporophore production of *Pleurotus* spp. (Jandaik 1974). Zedrazil (1980) has studied the effect of supplementing straw substrates with ammonium nitrate, soybean meal or alfalfa meal on the fructification and yield coefficient. Organic supplements affected not only yield raise but also raise in protein content and higher yield coefficient. Addition of organic supplements in the form of horse gram powder, cotton seed powder yeast mud, ground nut cake or rice bran also showed enhance yields coupled with increase in the protein content of fruit bodies of the several nitrogen sources tried as supplements to the rice straw substrate during the growth and fructification of *Pleurotus* species, oil seed cakes proved ideal in increasing the mushroom yield and its protein content (Rajaratnam *et al.* 1986, Bano *et al.* 1993). Facoya *et al.* (2014) cultivated *Pleurotus sajor-caju* on *Pycnanthus angolensis* sawdust supplemented with 0, 5, 10, 15, and 20% palmkernel cake, oil palm fibre, rice bran, wheat chaff, and corn cobs. Moda *et al.* (2005) cultivated the *Pleurotus sajor-caju* on sugarcane bagasses with supplementation with corn grits and mineral solution. Gupta *et al.* (2016) were used residual

biogas slurries of mahua cake and cow dung for supplementation in *Pleurotus sajor caju* for enhancement of nutritional contents. Sao and Deshmukh (2018) done the comparative study of natural (wheat flour, rice flour, soya flour, maize flour with Cotton seed cake) and chemical supplements (Lactose, ZnSO₄, MgSO₄, FeSO₄ with Peptone) for yield and biological efficiency of *Pleurotus sajor caju*. The survival and multiplication of mushrooms is related to a number of factors, which may act individually or have interactive effects among them. The combination of the best air temperature, moisture, nutrient conditions as well as other variables, provides a synergistic effect optimizing the production of mushrooms, with a consequent loss and cost reduction (Bulletini *et al.* 2019).

In order to select cheaper and better amendment of bio

efficiency of *P. sajor caju* an experiment was conducted in completely randomize design. As it is clearly evident from the result that supplement of *Embelica officinalis* not only reduced the spawn run time but also increases yield of *Pleurotus sajor-caju* in all three test lignocelluloses conventional substrate (wheat straw, paddy straw and soybean straw) and non-conventional substrates (domestic waste, fruit waste and used tea leaves). Supplementation of 3% concentration of *Embelica officinalis* gave best result in all three conventional and non-conventional substrate tested. These supplements were selected purposely because both have some nutritional values, easily available and their application can be exploited in production of potential food for men. So, they have used with positive hope of effectiveness.

LITERATURE CITED

- Baysal E, Peker H, Yalinkili M K and Temiz A. 2003. Cultivation of oyster mushroom on waste paper with some added supplementary materials. *Bioresource Technology* **89**(1): 95-97.
- Bano Z, Shashirekha M N and Rajarathnam S. 1993. Improvement of the bioconversion and biotransformation efficiencies of the oyster mushroom (*Pleurotus sajor caju*) by supplementation of its rice straw with oil seed cakes. *Enzymes and Microbial Technology* **15**: 985-989.
- Bahukhandi D. 1990. Effect of various treatments on paddy straw on yield of some cultivated species of *Pleurotus*. *Indian Phytopathology* **43**: 471-472.
- Bellettini M B, Assumpc F, Helayne F, Maieves A, Teixeira G La, vila a S A, Hornung P S, Maccari A and Ribani R H. 2019. Factors affecting mushroom *Pleurotus* spp. *Saudi Journal of Biological Sciences* **26**: 633-646.
- Bisaria R, Madan M and Bisaria V S. 1987. Biological efficiency and nutritive value of *Pleurotus sajor-caju* cultivated on different agro-wastes. *Bio Waste* **19**: 239-255.
- Bonatti M, Karnopp P, Soares H M and Furlan A S. 2003. Estudo da composicao de cogumelos das especies *Pleurotus ostreatus* and *Pleurotus sajor-caju* cultivos em palha de bananeira. *Rev Saude Amb* **4**: 31-35.
- Carrasco J, Zied D C, Pardo J E, Preston G M and Pardo-Gimenez A. 2018. Supplementation in mushroom crops and its impact on yield and quality. *AMB Expr* **8**: 146.
- Chang S T and Miles P G. 1989. *Edible Mushrooms and their Cultivation*. Florida: CRC Press.
- Chang S T and Miles P G. 2004. *Mushrooms-Cultivation, Nutritional Value, Medicinal Effect and Environmental Impact*. Second Edition. CRC Press. pp 2-4.
- Dias E S, Koshikumo E M S, Schwan R F and Silva R. 2003. Cultivation of the mushroom *Pleurotus sajor-caju* in different agricultural residues. *Ciën. Agrotec* **27**: 1363-1369.
- Dehariya P, Chaubey A and Vyas D. 2011. Effect of proteinaceous substrate supplementation on yield of *Pleurotus sajor-caju*. *Indian Phytopathology* **64**(3): 291-295.
- Fakoya S, Adejumo A F and Akinyele J B. 2014. Effect of the use of *Pycnanthus angolensis* and different supplements on yields and on the proximate composition of *Pleurotus sajor-caju*. *Journal of Mycology*. <http://dx.doi.org/10.1155/2014/642807>.
- Flack R. 1917. Uber die walkulter des austernpilzes (*Agaricus ostreatus*) out laubholzstubben. *Z. forst-sagdwes* **49**: 159-165.
- Gupta A and Raina P K. 2008. Effect of supplements on the yield of some high adaptive *Pleurotus species* in sub-tropics of Jammu. *Mushroom Research* **17**: 13-17.
- Gupta A, Sharma S, Kumar A, Alam P and Ahmad P. 2016. Enhancing nutritional contents of *Lentinus sajor-caju* using residual biogas slurry waste of detoxified mahua cake mixed with wheat straw. *Frontier Microbiology* **7**: 1529. doi: 10.3389/fmicb.2016.01529.
- Gunasegaran K and Graham K M. 1987. Effect of organic additives on yield of the Phoenix mushroom grown on cellulose waste. *Mushroom Journal of Tropics* **7**: 101-106.
- Gurjar K L and Doshi A. 1995. Effect of substrate supplements on fruit bodies production of *Pleurotus cornucopiae* (Paul ex Pers.) Rolland. *Mushroom Information* **10/12**: 12-23.
- Jain A K. 2005. Studies on cultivation technology of *Pleurotus* species and their marketing potential in Sagar region. *Ph. D. Thesis*.
- Jain A K and Vyas D. 2005. Supplementation of soybean choker: Enhances the growth and yield of *P. sajor-caju* grown in lignocellulosic waste. *Journal of Basic and Applied Mycology* **3/4**: 88-90.
- Jandaik C L. 1974. Artificial cultivation of *Pleurotus sajor-caju* mushroom. *Mushroom Journal* **22**: 405.

- Jeznabadi E K, Jafarpour M and Eghbalsaied S. 2016. King oyster mushroom production using various sources of agricultural wastes in Iran. *International Journal of Recycling and Organic Waste Agriculture* **5**: 17-24.
- Kalmis E and Sargin S. 2004. Cultivation of two *Pleurotus* sp. on wheat straw substrates containing olive mill waste water. *Int. Biodeter. Biode-grad* **53**: 43-47.
- Mishra A D and Singh C S. 2007. Effect of supplementation of wheat straw with resinous gums on the growth and production of *Pleurotus sajor-caju*. *Journal of Basic Applied Mycology* **6**: 3-4.
- Moonmoon M, Jahan N, Shelly M J, Khan M A, Uddin M N, Hossain K, Tania M and Ahmed S. 2011. Effects of different levels of wheat bran, rice bran and maize powder supplementation with saw dust on the production of shiitake mushroom (*Lentinus edodes* (Berk.) Singer). *Saudi Journal of Biological Science* **18**(4): 323-328.
- Moda E M, Horii J and Spoto M H F. 2005. Edible mushroom *Pleurotus sajor-caju* production on washed and supplemented sugarcane bagasse. *Science Agriculture* **62**: 127-132.
- Naraian R, Sahu R K, Kumar S, Garg S K, Singh C S and Kanaujia R S. 2009. Influence of different nitrogen rich supplements during cultivation of *Pleurotus florida* on corn cob substrate. *Environmentalist* **29**: 1-7.
- Naraian R, Srivastava J and Garg S K. 2011. Influence of dairy spent wash (DSW) on different cultivation phases and yield response of two *Pleurotus* mushrooms. *Annual of Microbiology* **61**: 853-862.
- Naraian R, Singh M P and Ram S. 2016. Supplementation of basal substrate to boost up substrate strength and oyster mushroom yield: An overview of substrates and supplements. *International Journal of Current Microbiology and Applied Science* **5**(5): 543-553.
- Peng J T, Lee C M and Tsai Y F. 2000. Effect of rice bran on the production of different king oyster mushroom strains during bottle cultivation. *Journal of Agricultural Research China* **49**(3): 60-67.
- Pedra W N and Marino R H. 2006. Cultivo axênico de *Pleurotus* spp. em serragem da casca de coco (*Cocos nucifera* Linn.) suplementada com farelo de arroz e/ou de trigo. *Arquivos do Instituto Biológico, São Paulo* **73**(2): 219-225.
- Pokhrel C P, Kalyan N, Budathoki U and Yadav R P. 2013. Cultivation of *Pleurotus sajor-caju* using different agricultural residues. *International Journal of Agricultural Policy and Research* **1**(2): 019-023.
- Patel Y, Naraian R and Singh V K. 2012. Medicinal Properties of *Pleurotus spp* (oyster mushroom): A review. *World Journal of Fungal and Plant Biology* **3**(1): 01-12.
- Rajarithnam S, Bano Z and Patwardhan M V. 1986. Nutrition of the mushroom *Pleurotus flabellatus* during its growth on paddy straw substrate. *Journal of Horticultural Science* **61**(2): 223-232, DOI: [10.1080/14620316.1986.11515694](https://doi.org/10.1080/14620316.1986.11515694)
- Royse D J and Schisler L C. 1986. Effect of benomyl application and spawn rate supplementation on yield and size of selected genotypes of *Pleurotus* spp. In: Proceedings of International Scientific and Technical Aspects of Cultivating Edible Fungi. Penn State University, Elsevier Science Publishers, Amsterdam. pp 109-115.
- Sao S and Deshmukh Y. 2018. Comparative study of natural and chemical supplementation effect on the yield and biological efficiency of *Pleurotus sajor-caju*. *International Journal of Advances in Science Engineering and Technology* **6**(1): 27-32.
- Silva E G, Dias E S, Siqueira F G and Schwan R F. 2007. Analise química de corpos de frutificacao de *Pleurotus sajor-caju* cultivado em diferentes cocentacoes de nitrogenio. *Cienc. Technology Alim.* **27**: 72-75.
- Shashirekha M N, Rajarithnam S and Bano Z. 2002. Enhancement of bioconversion efficiency and chemistry of the mushroom, *Pleurotus sajor-caju* (Berk and Br.) Sacc. Produced on spent rice straw substrate, supplemented with oil seedcakes. *Food Chemistry* **76**: 27-31. doi:10.1016/S0308-8146(01)00244-8
- Vijay B and Upadhyay R C. 1989. Chicken manure as a new nitrogen supplement in oyster mushroom cultivation. *Indian Journal of Mycology and Plant Pathology* **19**: 297-298.
- Zhang R, Xiujin L and Fadel J G. 2002. Oyster mushroom cultivation with rice and wheat straw. *Bioresource Technology* **82**: 277-284.
- Zadrazil F. 1980. Influence of ammonia nitrate and organic supplements on the yield of *Pleurotus sajor-caju* (Fr) Singer. *European Journal of Applied Microbiology and Biotechnology* **9**: 31-35.