

## Effect of *Ipomoea biloba* Leaves on Cardioprotective Activity against the Isoproterenol Induced Myocardial Infarction in Rats

R. Vallipriya<sup>\*1</sup> and M. Shabana Begum<sup>2</sup>

Received: 05 Nov 2020 | Revised accepted: 24 Jan 2021 | Published online: 09 Feb 2021  
© CARAS (Centre for Advanced Research in Agricultural Sciences) 2021

### ABSTRACT

The main aim of this investigation is to investigate the cardioprotective effect of ethanolic extract of *Ipomoea biloba* on isoproterenol induced myocardial infarction in rats. The rats were divided into five groups of six animals each. Group I served as a normal control; Group II rats were administered isoproterenol (20mg/kg) through sub cutaneous injection to induce the myocardial infarction. Group III and IV were pretreated with ethanolic extract of *Ipomoea biloba* leaf (100mg/kg, 200mg/kg respectively) before to the subcutaneous injection of isoproterenol (20mg/kg, b.w) for 2 consecutive days. Group V received ethanolic extract of *Ipomoea biloba* (200mg/kg b.w) alone for 28 days excluding the treatment procedures. After the completion of experimental period, the levels of cholesterol, triglycerides, phospholipids, and lipoproteins were estimated. The myocardial damage was assessed by quantifying the serum levels of cardiac marker enzymes (AST, ALT, LDH, CK, and TP) uric acid, and ceruloplasmin. The results of this experiment revealed that the Isoproterenol induced rats showed a significant increase in the levels of triglycerides, total cholesterol and phospholipids in both serum and heart homogenate. A rise in the levels of LDL, VLDL with significant decrease in the level of HDL was also observed in the serum of isoproterenol-intoxicated rats. Interestingly our findings were proved that the treatment with the ethanolic extract of *I. biloba* leaf was showed the noticeable decrease in the lipid profiles and myocardial marker enzymes in the experimental rats. The extract treatment also prevented the elevation of cholesterol, triglycerides and phospholipids in myocardial infarction induced rats. Hence, the findings of this work was clearly proved that the *I. biloba* leaf was appreciably prevented the isoproterenol induced myocardial infarction in rats.

**Key words:** Cardioprotective activity, *Ipomoea biloba*, Isoproterenol, Myocardial infarction

Myocardial infarction is a clinical disorder which is characterized by insufficient blood (without oxygen) supply from the coronary arteries and it results in damage or destroy of heart muscle [1-3]. The numerous previous studies showed that the oxidative stress produced by the generation of reactive oxygen species (ROS) plays a major role in the development of Myocardial infarction. It may sometimes damage cell membrane and metabolic alterations leads to cardiac dysfunction and damage [4-5]. In developing countries due to lifestyle changes myocardial infarction makes an increase in mortality and morbidity rate [6-7]. Isoproterenol (ISO) is a synthetic catecholamine and  $\beta$ -adrenergic agonist; on oxidation due to its free radicals, it produces stress in the myocardium and cause damage to the myocardial membrane [8]. Medicinal plants were demonstrated to play an important role in the management of various diseases in humans; including cardiovascular diseases [9]. *Ipomoea* is the largest genus in the flowering plant family Convolvulaceae, with more than 500 species. Some of the species of *Ipomoea* are

frequently used in Indian traditional medicine. The *Ipomoea biloba* is an important medicinal plant India with the numerous pharmacological activities. Many previous reports highlighted the biological activities of *Ipomoea biloba* plant [10-11]. However, still now there were no any scientific reports to claim the cardioprotective actions of *Ipomoea biloba* leaves. Hence this present investigation was aimed to investigate the cardioprotective activity of ethanolic leaf extract of *Ipomoea biloba* leaves against the ISO induced myocardial infarction in rats.

### MATERIALS AND METHODS

#### Collection of plant

The fresh and matured leaves of *Ipomoea biloba* were collected from Tamil Nadu coastal area. Then the collected leaves were washed completely with distilled water add the same was processed and dehydrated under the shady cabin.

#### Preparation of plant material

The shade dried plant leaves were powdered with an electrical blender and then the 20gm of *I. biloba* leaf powder was mixed with 100 ml of ethanol. Then it was heat macerated at the 85°C for 30 minutes and then the suspension was filtered by using the Whatman filter paper. Then the resulted

\*R. Vallipriya  
vallipriya2008@gmail.com

<sup>1,2</sup>Department of Biochemistry, Muthayammal College of Arts and Science, Rasipuram - 637 408, Namakkal District, Tamil Nadu, India

plant extract was powdered by vacuum evaporation process and finally the powder was used for the further investigations.

#### Experimental animals

The adult male Wistar albino rats were used for the study and the same was collected from the Institutional animal house. All the experimental animals were maintained under the standard laboratory conditions. All experimental animals were acclimatized for 7 days in prior to the starting of experiments and during that period animals were fed with standard pelleted rat chow and water *ad libitum*.

#### Experimental design

The rats were divided into five groups of six animals each. Group I: Served as a normal control. Group II: Rats were administered isoproterenol (20mg/kg) by the subcutaneous injection to induce the myocardial infarction. Group III and IV: Rats were pretreated with the ethanolic extract of *Ipomea biloba* leaf extract (100 and 200mg/kg, respectively) for a period of 28 days subsequently to the subcutaneous injection of isoproterenol (20mg/kg, b.w) for 2 consecutive days. Group V: Rats were received the ethanolic extract of *I. biloba* (200mg/kg b.w) alone for 28 days without any experimental treatments. After the experimental period, blood and heart tissue samples were collected and serum was separated and used for estimation of cholesterol, triglycerides, phospholipids, lipoproteins, myocardial marker enzymes, and other biochemical studies.

#### Preparation of heart tissue homogenate

After the completion of experiments, all the experimental animals were anesthetized by chloroform administration and sacrificed by cervical decapitation. After the animal scarification, the heart tissues were excised washed thoroughly in ice-cold phosphate buffered saline to remove the excess blood. Ten percent of homogenate was prepared in 0.1M Tris HCl buffer (pH-7.4). Then the homogenate was centrifuged at 6000 rpm for 20 min at 4°C and the supernatant was used for the further biochemical assays.

#### Estimation of cholesterol

The total cholesterol in both serum and heart tissue sample were estimated according to the previous method described [12]. The 0.1ml of the samples was added to the 4.9 ml of ferric chloride precipitation reagent and then centrifuged for 6 minutes at 7000rpm and then the resulting supernatant was collected. The 2.5 ml of supernatant was added to the 2.5 ml of ferric chloride diluting reagent and along with the 4 ml of concentrated sulphuric acid. The reaction mixture was incubated for 10 minutes at room temperature and finally optical density was measured at 560 nm. The cholesterol content was expressed as mg/dl of serum.

#### Determination of triglyceride

The triglycerides level in the serum and heart tissue homogenate was estimated by the method [13]. Briefly, the 1.0 ml of isopropanol was added to 0.1 ml of each sample and the 4g of alumina was added and then the suspension was shaken well for 15 mins. Then this suspension was centrifuged at 6000 rpm for 7 minutes then 2.0ml of the supernatant was placed to the properly labelled vials. Then the tubes were located in the water bath at 70°C for 20 min for saponification after adding 0.6 ml of the saponification reagent. Finally, the suspension was cooled to room temperature then 1.0 ml of sodium metaperiodate was added subsequently to the 0.5 ml of

acetyl acetone reagent. The contents were cooled and the absorbance was taken at 430nm.

#### Determination of phospholipids

The previously described method by [14] was employed to determine the phospholipids in the samples. For this, the 0.1 ml of each sample was treated with the 0.2 ml of perchloric acid and this digestion was continued till the reaction solution becomes colourless. After that, the liberated phosphorus was estimated adding 4.3 ml of H<sub>2</sub>O to the digested sample and 0.5 ml of ammonium molybdate. Then the tubes were shaken well and maintained for 20 minutes incubation. Finally, the developed blue colour in the reaction solution was determined at 620 nm.

#### Assay of lipoproteins level

The levels of lipoproteins such as, high density lipoprotein (HDL), low density lipoprotein (LDL), very low density lipoprotein (VLDL) in the serum as well as heart tissue homogenate of the experimental animals were estimated according to the previous described method [15]. Briefly, the 1 ml serum and heart tissue homogenate were added separately to the 0.1 ml of phosphotungstate reagent and 50µl of magnesium chloride reagent. The reaction content was centrifuged at optimum temperature for 30 minutes at 3000 rpm. Then the 0.1 ml of resulted supernatant was added to the 4.9 ml of ferric chloride precipitating reagent, then it was mixed well and again centrifuged at 3000rpm for 10 minutes. After that, 2.5 ml of resulting supernatant was taken and mixed with the 2.5 ml of diluting reagent and 4 ml of concentrated sulphuric acid by mixing thoroughly. Subsequently, the blank was also maintained for the results interpretations. The colour developed was read at 560nm.

#### Biochemical analysis

Cardiotoxicity was assessed by quantifying the serum levels of aspartate transaminase (AST), alanine transaminase (ALT), lactate dehydrogenase (LDH), creatine kinase (CK), total protein (TP), uric acid, and the ceruloplasmin, by using the commercially procured assay kits (Sigma aldrich, USA).

#### Statistical analysis

All conclusions happen to be demonstrated as Mean  $\pm$  SD. in each group for six animals. All the compiled data were statistically examined utilizing SPSS10 software. Theory Hypothesis test approaches included one-way variance analysis (ANOVA) followed by the least significant difference (LSD) test. The significance level at alpha=0.05 was considered to statistical significance.

## RESULTS AND DISCUSSION

#### Effects of ethanolic extract of *Ipomea biloba* leaves on cholesterol, triglycerides, and phospholipids level in the experimental rats

The myocardial infarction was induced to the experimental animals by the injection of ISO by subcutaneously to examine the therapeutic potential of ethanolic leaf extract of *I. biloba*. The subcutaneous injection ISO to the rats (Group II) showed the drastic increase in the cholesterol and triglycerides level in the serum as well as heart tissue homogenate when compared to the control rats (Group I). The difference in the levels of cholesterol and triglycerides in the control and myocardial infarction induced rats were contrasts each other. Interestingly, the treatment with the

ethnolic extract of *Ipomea biloba* leaves (100 and 200mg/kg) to the myocardial infarction induced rats showed the noticeably decreased level of the both cholesterol and triglycerides (Fig 1). The extract alone treated animals were showed completely similar kind of results to the control. There were no any significant variations between control and plant extract treated experimental animals.




Fig 1 Effects of ethanolic extract of *Ipomea biloba* leaves on the levels of cholesterol, triglycerides, and phospholipids in the experimental rats

*Effects of ethanolic extract of Ipomea biloba leaves on lipoproteins level in the experimental rats*




Fig 2 Effect of ethanolic extract of *Ipomea biloba* leaf on the lipoproteins level in the experimental rats





Fig 3 Effect of ethanolic extract of *Ipomea biloba* leaf on the myocardial marker enzymes level in the experimental rats

*Effects of ethanolic extract of Ipomea biloba leaves on marker enzymes level in the experimental rats*

As depicted in the (Fig 3), the levels of aspartate transaminase (AST), alanine transaminase (ALT), lactate dehydrogenase (LDH), and creatine kinase (CK) was drastically elevated and also the reduced level of total proteins were noted in the serum and heart tissue homogenate of the myocardial infarction stimulated experimental animals when compared to the untreated control animals. This severs

In this present study, ISO induced myocardial infarction in rats were demonstrated the significantly elevated levels of low-density lipoprotein (LDL), very low-density lipoprotein (VLDL), and decreased the high-density lipoprotein (HDL) in the serum as well as heart tissue homogenate of the myocardial induced experimental rats when compared to the control group. This result showed the severe effects of ISO induced myocardial infarction in rat model. Whereas, the 100 and 200mg/kg of ethnolic extract of *Ipomea biloba* leaves (100 and 200mg/kg) treatment to the myocardial infarction induced rats demonstrated the appreciably reduced levels low density lipoprotein (LDL), very low-density lipoprotein (VLDL), and amazingly increased level of high-density lipoprotein (HDL) when compared to the myocardial induced experimental rats (Fig 2). This result was proved the beneficial activity of ethanolic extract of *Ipomea biloba* leaves against the cardio-protection. There were no any significant variations between control and plant extract treated experimental animals. The extract alone treated animals were showed completely similar kind of results to the control.



reduction in the myocardial markers enzymes in the experimental rats were revealed the serious myocardial infarction in the experimental animals. Interestingly the treatment with the 100 and 200mg/kg of ethnolic extract of *Ipomea biloba* leaves (Group III and IV) to the myocardial infarction induced rats showed the amazingly reduced levels of myocardial marker enzymes such as, AST, ALT, LDH and CK in the serum as well as heart tissue homogenate when compared to the myocardial infarction induced experimental

rats. The extract treatment also revealed the significantly increased levels of total proteins in the serum and heart tissue homogenate of the myocardial infarction induced experimental rats (Fig 3). The extract alone treated animals did not show any alterations in the myocardial marker enzymes and it was quite similar to results of the control. There were no any significant variations were noted between control and plant extract treated experimental animals (Fig 3).

#### *Effects of ethanolic extract of Ipomea biloba leaves on the uric acid and ceruloplasmin level in the experimental rats*

Data depicted in (Fig 4) revealed that the level of uric acid was harshly increased and the level of ceruloplasmin protein was drastically decreased in the serum of myocardial infarction induced (Group II) experimental animals when compared to the untreated normal control animals (Group I). The severe increase of the uric acid in the serum of myocardial infarction induced rats revealing the declined activity and kidneys. On the other hand, the treatment with the ethanolic extract of the *Ipomea biloba* leaves (100 and 200mg/kg) to the myocardial infarction induced experimental animals (Group III and IV) were showed the significantly reduced level of uric acid and increased level of ceruloplasmin in the serum of myocardial infarction induced experimental animals than the diseased animals (Group II). The extract treatment appreciably regained the ceruloplasmin level and reduced the uric acid level in the serum of experimental animals (Fig 4). There were no any significant variations were noted between control and plant extract treated experimental animals. The extract alone treated animals did not show any alterations in the uric acid and ceruloplasmin levels in the serum of experimental animals.




Fig 4 Effect of ethanolic extract of *Ipomoea biloba* leaf on uric acid and ceruloplasmin levels in the serum of experimental rats

Myocardial infarction is characterized by altered cardiac function due to imbalance between oxygen supply and demand. Oxidative stress induced by reactive oxygen species (ROS) plays a major role. ROS mediate various signaling pathways that results into vascular inflammation and atherogenesis. Oxidative stress by the reactive oxygen species is mainly responsible for myocardial infarction. Isoproterenol is responsible for the free radical formation which may cause cellular cholesterol accumulation by increasing the cholesterol accumulation and decrease in its utilization [16-17]. In this present study, the findings were exactly similar to this statement. The results of this study showed that the isoproterenol injection to the experimental animals were demonstrated the gradual increase levels of cholesterol, triglycerides and phospholipids in the serum as well as heart tissues of the myocardial infarction induced experimental

animals. This causes the severe damage to the myocardial membrane. Interestingly, the treatment with the ethanolic extract of *Ipomea biloba* leaves to the myocardial infarction induced rats showed the noticeably decreased level of the both cholesterol and triglycerides (Fig 1). There were no any significant variations between control and plant extract treated experimental animals.

The Myocardium contains the abundant of cardiac enzymes such as aspartate aminotransferase (AST), alanine aminotransferase (ALT), lactate dehydrogenase (LDH), and creatine kinase (CK). These enzymes are some of the ideal markers for the indication of myocardial infarction. The drastic elevation of these marker enzymes levels in the serum, denoting the severe injury to the myocardium. The injury to the cardio tissue may release these marker enzymes to the serum. In this study, we also noted the exact outcomes. The isoproterenol induced animals (Group II) showed the severely elevated levels of these marker enzymes such as AST, ALT, LDH, and CK in the both serum and heart tissues. This elevation in these marker enzymes may lead to the administration of isoproterenol, which may lead to the insufficient supply of oxygen to the cardio cells that further leads to cell cellular damage and/or tissue necrosis. This phenomenon allows the cardiac marker enzymes to leak out in to the blood stream [18]. The findings of our present study were revealed that the pre-treatment with ethanolic extract of *Ipomoea biloba* leaves were significantly decreases the levels of cardio marker enzymes in the serum of experimental animals and showed the appreciable cardio-protection.

A few previous research reports were confirmed that the uric acid is prime indicator of the kidney function that directly associated with the heart diseases. In the case of myocardial infarction, the uric acid level in the serum was regarded as the foremost risk factor. The uric acid is usually produced from the xanthine. During the hypoxic conditions, the tissues are disturbed due to the increased oxidation of essential-SH-groups of the enzyme xanthine dehydrogenase and it was converted to xanthine oxidase. Xanthine oxidase enzymes normally catalyze the conversion of hypoxanthine to xanthine, uric acid and superoxide. The uric acid can also stimulate the granulocytes adherence to the endothelium and peroxide and superoxide free radical liberation [19-20]. These free radicals liberation eventually produces the increased stress to the myocardium and causes cellular damage and/or cell death. In this investigation, we also noted the drastically increased level of the uric acid in the serum of myocardial infarction induced experimental animals than the untreated normal control animals. Interestingly, the treatment with the ethanolic extract of the *Ipomea biloba* leaves to the myocardial infarction induced experimental animals were showed the appreciably reduced level of uric acid in the serum of myocardial infarction induced experimental animals (Fig 4). There were no any significant variations were noted between control and plant extract treated experimental animals.

The ceruloplasmin is an extracellular antioxidant and acute phase protein. Some previous findings were demonstrated that the ceruloplasmin protects the intima against free radical stimulated tissue injury. The antioxidant property of ceruloplasmin is through its ferroxidase activity catalyzing the oxidation of  $\text{Fe}^{2+}$  to  $\text{Fe}^{3+}$ . Thus, it inhibits ferrous ion stimulated lipid peroxidation. Since it has both ferroxidase and copper binding capacity, it is used to neutralize the excess amount of free radicals in the body [21-22]. With the above note, in this study we have also observed

the significantly decreased levels of ceruloplasmin in the serum of ISO induced myocardial infarction rats. The findings of this present study were revealed that the pre-treatment with ethanolic extract of *Ipomoea biloba* leaves were noticeably augmented the ceruloplasmin protein level in the serum of experimental animals.

## CONCLUSIONS

In conclusion, the novel findings of this present investigation were proved that the ethanolic extract of *Ipomoea biloba* leaves were showed the appreciable cardio

protection against the ISO-induced myocardial infarction in experimental rats. The ethanolic extract treatment noticeably reduced the cholesterol, triglycerides, myocardial marker enzymes, and low-density lipoproteins levels in both serums as well as heart tissue. These results were proved the cardio-protective action of the ethanolic extract of *Ipomoea biloba* leaves. Hence, it was concluded that the *Ipomoea biloba* leaves may play a significant role in the development of novel cardio-protective drugs in future. However, the additional researches were still needed in future to elucidate the exact curative mechanism of the *Ipomoea biloba* leaves against the myocardial infarction.

## LITERATURE CITED

1. Abi-Beaulah G, Mohamed SA, Sivakumar V, Jaya SR. 2014 Cardioprotective activity of methanolic extract of *Croton sparciflorus* on isoproterenol induced myocardial infarction wistar albino rats. *Journal of Medicinal Plants* 2(6): 01-8.
2. Goyal SN, Sharma C, Mahajan UB, Patil CR, Agrawal YO, Kumari S, Arya DS, Ojha S. 2015. Protective effects of cardamom in isoproterenol-induced myocardial infarction in rats. *International Journal of Molecular Sciences* 16(11): 27457-69.
3. Hayat Malik MN, Bashir S, Khan IU, Karim S, Mushtaq MN, Khan HU, Rashid M, Naz H, Samreen S. 2015. Cardiotonic and vasoconstriction effects of aqueous methanolic extract of *Paspalum flavidum* L. *Pakistan Journal of Pharmaceutical Sciences* 28(2): 421-424.
4. Ojha SK, Bharti S, Joshi S, Kumari S, Arya DS. 2012. Protective effect of hydroalcoholic extract of *Andrographis paniculata* on ischaemia-reperfusion induced myocardial injury in rats. *The Indian Journal of Medical Research* 135(3): 414.
5. Wang Y, Zhang ZZ, Wu Y, Zhan J, He XH, Wang YL. 2013. Honokiol protects rat hearts against myocardial ischemia reperfusion injury by reducing oxidative stress and inflammation. *Experimental and Therapeutic Medicine* 5(1): 315-319.
6. Nandave M, Ojha SK, Kumari S, Nag TC, Mehra R, Narang R, Arya DS. 2013. Cardioprotective effect of root extract of *Picrorhiza kurroa* (Royle Ex Benth) against isoproterenol-induced cardiotoxicity in rats. *Indian Journal of Experimental Biology* 51: 694-701.
7. Boudina S, Laclau MN, Tariosse L, Daret D, Gouverneur G, Bonoron-Adèle S, Saks VA, Dos Santos P. 2002. Alteration of mitochondrial function in a model of chronic ischemia *in vivo* in rat heart. *American Journal of Physiology-Heart and Circulatory Physiology* 282(3): 821-831.
8. Mohanty IR, Arya DS, Gupta SK. 2009. Dietary *Curcuma longa* protects myocardium against isoproterenol induced hemodynamic, biochemical and histopathological alternations in rats. *International Journal of Applied Research in Natural Products* 1(4): 19-28.
9. Barman NR, Nandy S, Datta R, Kar PK. 2013 Sep; Cardioprotective effect of ethanolic extract of *Urtica parviflora* Roxb. against isoproterenol induced myocardial infarction in rats. *Indian Journal of Pharmacology* 45(5): 513.
10. Venkataraman ND, Atlee WC, Prabhu TP. 2013. Anti-inflammatory potential of ethanolic extracts from aerial parts of *Ipomoea pes-caprae* (L.) R. Br using cotton pellet induced granuloma model. *Journal of Applied Pharmaceutical Science* 3(7): 61.
11. Matunog VE, Bajo LM. 2013. Phytochemical Screening and Antioxidative Potentials of “Beach Morning Glory” *Ipomoea pes-caprae* (Linn.) Roth Leaves Extract. *Journal of Multidisciplinary Studies* 1(1): 1-18.
12. Zak B, Dickenman RC, White EG, Burnett H, Cherney PJ. 1954. Rapid estimation of free and total cholesterol. *American Journal of Clinical Pathology* 24(11): 1307-1315.
13. Foster LB, Dunn RT. 1973. Stable reagents for determination of serum triglycerides by a colorimetric Hantzsch condensation method. *Clinical Chemistry* 19(3): 338-340.
14. Bartlett GR. 1959 Phosphorus assay in column chromatography. *Journal of Biological Chemistry* 234: 466-448.
15. Friedewald WT, Levy RI, Fredrickson DS. 1972. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clinical Chemistry* 18(6): 499-502.
16. Khan G, Haque SE, Anwer T, Ahsan MN, Safhi MM, Alam MF. 2014. Cardioprotective effect of green tea extract on doxorubicin-induced cardiotoxicity in rats. *Acta Pol Pharm.* 71(5): 861-868.
17. Shi M, He W, Liu Y, Li X, Yang S, Xu Q. 2013. Protective effect of total phenylethanoid glycosides from *Monochasma savatieri* Franch on myocardial ischemia injury. *Phytomedicine* 20(14): 1251-1255.
18. Prabha SP, Nitha A, Ansil PN, Latha MS. 2014. Cardioprotective effect of methanolic extract of *Gardenia gummifera* Linn. F. on isoproterenol induced myocardial infarction in rats. *Int. Jr. of Pharm Sci. Res.* 5: 3817-3828.
19. Rameshrad M, Vaez H, Toutounchi NS, Fathiazad F, Garjani A. 2014. Effect of methanolic extract of *marrubium crassidens* boiss on ischemia/reperfusion induced arrhythmias and infarct size in isolated rat heart. *Pharmaceutical Sciences* 20(3): 80-89.
20. Bae MH, Lee JH, Lee SH, Park SH, Yang DH, Park HS, Cho Y, Jun JE, Chae SC. 2011 Serum uric acid as an independent and incremental prognostic marker in addition to N-terminal pro-B-type natriuretic peptide in patients with acute myocardial infarction. *Circulation Journal* 75(6): 1440-1447.
21. Allahyari S, Delazar A, Najafi M. 2014. Evaluation of general toxicity, anti-oxidant activity and effects of *Ficus carica* leaves extract on ischemia/reperfusion injuries in isolated heart of rat. *Advanced Pharmaceutical Bulletin* 4(Suppl 2): 577.
22. Osaki S, Johnson DA, Frieden E. 1966. The possible significance of the ferrous oxidase activity of ceruloplasmin in normal human serum. *Journal of Biological Chemistry* 241(12): 2746-2751.